


SUPPLEMENTARY INFORMATION

Ribose 2'-*O*-methylation provides a molecular signature for MDA5-dependent distinction of self and non-self mRNA

Roland Züst¹, Luisa Cervantes-Barragan¹, Matthias Habjan¹, Reinhard Maier¹, Benjamin W. Neuman², John Ziebuhr^{3,4}, Kristy J. Szretter⁵, Susan C. Baker⁶, Winfried Barchet⁷, Michael S. Diamond⁵, Stuart G. Siddell⁸, Burkhard Ludewig^{1,9} and Volker Thiel^{1,9}


¹Institute of Immunobiology, Kantonal Hospital St. Gallen, St. Gallen, Switzerland. ²School of Biological Sciences, University of Reading, United Kingdom. ³Centre for Infection and Immunity, Queen's University Belfast, Belfast United Kingdom. ⁴Institute of Medical Virology, Justus Liebig University Giessen, Giessen, Germany. ⁵Departments of Medicine, Molecular Microbiology, and Pathology & Immunology, Washington University School of Medicine, St Louis, MO 63110, USA. ⁶Department of Microbiology and Immunology, Loyola University Stritch School of Medicine, Maywood, Illinois, USA. ⁷Institute for Clinical Chemistry and Pharmacology, University Hospital, University of Bonn, Germany ⁸Department of Cellular and Molecular Medicine, School of Medical and Veterinary Sciences, University of Bristol, Bristol, United Kingdom. ⁹Vetsuisse Faculty, University of Zürich, Switzerland.

Supplementary Figure 1



Supplementary Figure 1. Morphology, size and number of MHV-induced DMVs are not affected by 2'-O-MTase-deficiency. DMVs were analysed by electron microscopy of ultrathin sections of MHV-A59-, MHV-Y15A-, or MHV-D130A-infected (MOI=1) mouse L929 cells at 5 h p.i.. Virus-induced features including convoluted membranes (c), DMVs (*) and virions (v), as well as cellular features including the nucleus (n), Golgi (g), mitochondria (m) and ordered smooth endoplasmic reticulum (o) are indicated. Quantitative data describing DMV and virions visible in these micrographs are shown at lower right. The average of the longest and shortest visible diameter was taken as a measure of size. The ratio of the longest to the shortest diameter was taken as a measure of shape. The number of DMVs per cell includes only DMV-containing cells. Scale bar denotes 200 nm.

Supplementary Figure 2



Supplementary Figure 2. MHV 2'-O-MTase mutants induce increased IFN-I secretion. a,b)

Kinetics of IFN-I secretion induced by MHV 2'-O-MTase mutants was analyzed by bioassay for IFN-I. MΦs (1×10^6) derived from wild-type (a), or IFNAR-deficient (b) mice were infected (MOI=1) with MHV-A59, MHV-Y15A, or MHV-D130A and IFN-I in cell culture supernatants was determined at indicated time points p.i.. Results represent the mean \pm SEM of two independent experiments (n=4). Statistical analysis was performed using unpaired Student's t-test (**, p < 0.01).

Supplementary Table 1. Methylation of viral RNA in selected cytoplasmically-replicating virus families infecting multicellular eukaryotes.

Virus class or Order	Family	Genus	RrmJ-like Domain	Function	Reference	mRNA 5' Structure ^a	Reference	Notes
ssRNA(+) Viruses								
Nidovirales	Coronaviridae	All	nsp16	2OMT	47	^{7m} GpppA ^{2'Om} or ^{7m} GpppG ^{2'Om}	48,49	^b
Nidovirales	Roniviridae	Okavirus	MHV nsp16 homolog	MTh ^c	50	<i>not determined</i>		^d
Nidovirales	Arteriviridae	Arterivirus	nd ^e			^{7m} GpppA ^{2'Om} or ^{7m} GpppG ^{2'Om}	51	^f
Picornavirales	All	All	nd					
Tymovirales	All	All	nd			VPg and IRES	52	
-	Astroviridae	All	nd			^{7m} GpppG	53	^g
-	Barnaviridae	Barnavirus	nd			<i>not determined</i>		^f
-	Bromoviridae	All	nd			VPg	54	
-	Caliciviridae	All	nd			^{7m} GpppG	55	^g
-	Closteroviridae	All	nd			VPg	56	
-	Flaviviridae	Flavivirus	NS5	N7/2OMT	57	^{7m} GpppA ^{2'Om}	58	
-	Flaviviridae	Pestivirus	nd			IRES	59	
-	Flaviviridae	Hepacivirus	nd			IRES	60	
-	Hepeviridae	Hepevirus	nd			^{7m} Gppp	61	^{h,i}
-	Luteoviridae	All	nd			Uncapped	62	
-	Nodaviridae	All	nd			^{7m} GpppG	63	
-	Potyviridae	All	nd			VPg and IRES	64	
-	Tetraviridae	All	nd			<i>not determined</i>		^g
-	Togaviridae	All	nsp2	MTh ^j	65	^{7m} GpppA	66	^k
-	Tombusviridae	All	nd			^{7m} GpppA	67	
-	Virgaviridae	All	nd			^{7m} GpppG	68	^g
ssRNA(-) Viruses^l								
Mononegavirales	Rhabdoviridae	All	L	N7/2OMT	69	^{7m} GpppA ^{2'Om} or ^{7m} GpppN	70	
Mononegavirales	Filoviridae	All	L	MTh ^c	71	^{7m} Gppp	72	ⁱ
Mononegavirales	Paramyxoviridae	All	L	MTh ^c	73	^{7m} GpppG	74	
-	Arenaviridae	Arenavirus	nd			Methylated cap	75	^m
-	Bunyaviridae	All	nd			^{7m} Gppp	76	^{i,m}
dsRNA Viruses^m								
-	Birnaviridae	All	nd			VPg	77	
-	Reoviridae	All	$\lambda 2/\text{Cap}$	MTh ^c	78	^{7m} GpppN ^{2'Om}	79	^o
dsDNA Virusesⁿ								
-	Asfarviridae	Asfivirus	EP424R	MTh ^c	80	^{7m} GpppA ^{2'Om}	81	
-	Mimiviridae	Mimivirus	L511	MTh ^c	82	<i>not determined</i>		
-	Poxviridae	All	VV_vp39 homologs	2OMT	83	^{7m} Gppp ^{6m} A ^{2'Om}	84	^q
-	Phycodnaviridae	All	nd			<i>not determined</i>		^r

^aIncluding genomic RNA structure data for positive-sense single-stranded RNA viruses only.

^bCoronavirus nsp14 is an N7MT.

^cIndicates detection of one domain with 2OMT homology.

^dIt is unclear whether roniviruses also encode a homolog of the coronavirus nsp14 N7MT.⁴⁸

^eNot detected.

^f2OMT and N7MT homologs not detected.

^gOne domain has N7MT homology.^{65,85}

^hHepatitis E virus p110 has N7MT activity.⁸⁶

ⁱViral RNAs interact with ^{7m}G-cap binding antisera.

^jA homologous domain is present, but active site residues are not conserved.

^kOne domain of nsp1 has N7MT homology, and internal RNA methylation has been reported ^{65,66}.

^lOmitting bornavirus and nucleorhabdoviruses, which replicate in the nucleus.

^mThe cap structure of viral mRNA is acquired from host mRNA ^{87,88}.

ⁿOmitting the poorly characterized genomes of the Endornaviridae, Partitiviridae and Picobirnaviridae.

^oOmitting the poorly characterized genomes of Fijivirus Mimoreovirus and Dinohernavirus.

^pOmitting Iridoviridae which carry out most or all mRNA synthesis in the nucleus.

^qConserved poxvirus VV D1-like proteins have N7MT homology ⁸⁹.

^rHomologs of chlorella virus A103R have N7MT activity ⁹⁰

References for Supplementary Table 1

- ⁴⁷ Decroly, E. et al., Coronavirus nonstructural protein 16 is a cap-0 binding enzyme possessing (nucleoside-2' O)-methyltransferase activity. *J Virol* **82** (16), 8071 (2008).
- ⁴⁸ This study.
- ⁴⁹ van Vliet, A. L., Smits, S. L., Rottier, P. J., and de Groot, R. J., Discontinuous and non-discontinuous subgenomic RNA transcription in a nidovirus. *EMBO J* **21** (23), 6571 (2002); Lai, M. M. and Stohlman, S. A., Comparative analysis of RNA genomes of mouse hepatitis viruses. *J Virol* **38** (2), 661 (1981).
- ⁵⁰ Gorbalenya, A. E., Enjuanes, L., Ziebuhr, J., and Snijder, E. J., Nidovirales: evolving the largest RNA virus genome. *Virus Res* **117** (1), 17 (2006).
- ⁵¹ Sagripanti, J. L., Zandomeni, R. O., and Weinmann, R., The cap structure of simian hemorrhagic fever virion RNA. *Virology* **151** (1), 146 (1986).
- ⁵² Le Gall, O. et al., Picornavirales, a proposed order of positive-sense single-stranded RNA viruses with a pseudo-T = 3 virion architecture. *Arch Virol* **153** (4), 715 (2008).
- ⁵³ Briand, J. P., Keith, G., and Guilley, H., Nucleotide sequence at the 5' extremity of turnip yellow mosaic virus genome RNA. *Proc Natl Acad Sci U S A* **75** (7), 3168 (1978); Klein, C. et al., Physical and functional heterogeneity in TYMV RNA: evidence for the existence of an independent messenger coding for coat protein. *Nucleic Acids Res* **3** (11), 3043 (1976).
- ⁵⁴ Revill, P. A., Davidson, A. D., and Wright, P. J., Mushroom bacilliform virus RNA: the initiation of translation at the 5' end of the genome and identification of the VPg. *Virology* **249** (2), 231 (1998).
- ⁵⁵ Dasgupta, R., Shih, D. S., Saris, C., and KAESBERG, P., Nucleotide sequence of a viral RNA fragment that binds to eukaryotic ribosomes. *Nature* **256** (5519), 624 (1975); Pinck, L., The 5'-end groups of alfalfa mosaic virus RNAs are m7G5'ppp5'Gp. *FEBS Lett* **59** (1), 24 (1975).
- ⁵⁶ Schaffer, F. L., Ehresmann, D. W., Fretz, M. K., and Soergel, M. I., A protein, VPg, covalently linked to 36S calicivirus RNA. *J Gen Virol* **47** (1), 215 (1980).
- ⁵⁷ Ray, D. et al., West Nile virus 5'-cap structure is formed by sequential guanine N-7 and ribose 2'-O methylations by nonstructural protein 5. *J Virol* **80** (17), 8362 (2006).
- ⁵⁸ Cleaves, G. R. and Dubin, D. T., Methylation status of intracellular dengue type 2 40 S RNA. *Virology* **96** (1), 159 (1979).
- ⁵⁹ Poole, T. L. et al., Pestivirus translation initiation occurs by internal ribosome entry. *Virology* **206** (1), 750 (1995).
- ⁶⁰ Tsukiyama-Kohara, K., Iizuka, N., Kohara, M., and Nomoto, A., Internal ribosome entry site within hepatitis C virus RNA. *J Virol* **66** (3), 1476 (1992).
- ⁶¹ Kabrane-Lazizi, Y., Meng, X. J., Purcell, R. H., and Emerson, S. U., Evidence that the genomic RNA of hepatitis E virus is capped. *J Virol* **73** (10), 8848 (1999).
- ⁶² Allen, E., Wang, S., and Miller, W. A., Barley yellow dwarf virus RNA requires a cap-independent translation sequence because it lacks a 5' cap. *Virology* **253** (2), 139 (1999).
- ⁶³ Dasgupta, R. et al., Primary and secondary structure of black beetle virus RNA2, the genomic messenger for BBV coat protein precursor. *Nucleic Acids Res* **12** (18), 7215 (1984).
- ⁶⁴ Carrington, J. C. and Freed, D. D., Cap-independent enhancement of translation by a plant potyvirus 5' nontranslated region. *J Virol* **64** (4), 1590 (1990).
- ⁶⁵ Rozanov, M. N., Koonin, E. V., and Gorbalenya, A. E., Conservation of the putative methyltransferase domain: a hallmark of the 'Sindbis-like' supergroup of positive-strand RNA viruses. *J Gen Virol* **73** (Pt 8), 2129 (1992).
- ⁶⁶ Dubin, D. T. et al., Sindbis virus messenger RNA: the 5'-termini and methylated residues of 26 and 42 S RNA. *Virology* **77** (2), 457 (1977).
- ⁶⁷ Xiong, Z. and Lommel, S. A., The complete nucleotide sequence and genome organization of red clover necrotic mosaic virus RNA-1. *Virology* **171** (2), 543 (1989).
- ⁶⁸ Zimmern, D., The 5' end group of tobacco mosaic virus RNA is m7G5'ppp5'Gp. *Nucleic Acids Res* **2** (7), 1189 (1975); Haidar, M. A. and Hirth, L., 5'-terminal structure of tobacco rattle virus RNA: evidence for polarity of reconstitution. *Virology* **76** (1), 173 (1977).

- 69 Li, J., Wang, J. T., and Whelan, S. P., A unique strategy for mRNA cap methylation used by vesicular
70 stomatitis virus. *Proc Natl Acad Sci U S A* **103** (22), 8493 (2006).
- 70 Abraham, G., Rhodes, D. P., and Banerjee, A. K., The 5' terminal structure of the methylated mRNA
synthesized in vitro by vesicular stomatitis virus. *Cell* **5** (1), 51 (1975); Colonna, R. J. and Stone, H. O.,
71 Newcastle disease virus mRNA lacks 2'-O-methylated nucleotides. *Nature* **261** (5561), 611 (1976).
- 71 Feder, M., Pas, J., Wyrwicz, L. S., and Bujnicki, J. M., Molecular phylogenetics of the RrmJ/fibrillarin
superfamily of ribose 2'-O-methyltransferases. *Gene* **302** (1-2), 129 (2003).
- 72 Weik, M. et al., Ebola virus VP30-mediated transcription is regulated by RNA secondary structure
formation. *J Virol* **76** (17), 8532 (2002).
- 73 Bujnicki, J. M. and Rychlewski, L., In silico identification, structure prediction and phylogenetic analysis
of the 2'-O-ribose (cap 1) methyltransferase domain in the large structural protein of ssRNA negative-
strand viruses. *Protein Eng* **15** (2), 101 (2002).
- 74 Barik, S., The structure of the 5' terminal cap of the respiratory syncytial virus mRNA. *J Gen Virol* **74** (Pt
3), 485 (1993).
- 75 Carter, M. F., Biswal, N., and Rawls, W. E., Characterization of nucleic acid of pichinde virus. *J Virol* **11**
(1), 61 (1973).
- 76 Hacker, D., Rochat, S., and Kolakofsky, D., Anti-mRNAs in La Crosse bunyavirus-infected cells. *J Virol*
64 (10), 5051 (1990); Vialat, P. and Bouloy, M., Germiston virus transcriptase requires active 40S
ribosomal subunits and utilizes capped cellular RNAs. *J Virol* **66** (2), 685 (1992).
- 77 Persson, R. H. and Macdonald, R. D., Evidence that infectious pancreatic necrosis virus has a genome-
linked protein. *J Virol* **44** (2), 437 (1982).
- 78 Mao, Z. X. and Joklik, W. K., Isolation and enzymatic characterization of protein lambda 2, the reovirus
guanylyltransferase. *Virology* **185** (1), 377 (1991).
- 79 Both, G. W., Furuchi, Y., Muthukrishnan, S., and Shatkin, A. J., Ribosome binding to reovirus mRNA in
protein synthesis requires 5' terminal 7-methylguanosine. *Cell* **6** (2), 185 (1975).
- 80 Saha, N. and Shuman, S., Effects of alanine cluster mutations in the D12 subunit of vaccinia virus mRNA
(guanine-N7) methyltransferase. *Virology* **287** (1), 40 (2001).
- 81 Salas, M. L., Kuznar, J., and Vinuela, E., Polyadenylation, methylation, and capping of the RNA
synthesized in vitro by African swine fever virus. *Virology* **113** (2), 484 (1981).
- 82 Raoult, D. et al., The 1.2-megabase genome sequence of Mimivirus. *Science* **306** (5700), 1344 (2004).
- 83 Schnierle, B. S., Gershon, P. D., and Moss, B., Cap-specific mRNA (nucleoside-O2')-methyltransferase
and poly(A) polymerase stimulatory activities of vaccinia virus are mediated by a single protein. *Proc Natl
Acad Sci U S A* **89** (7), 2897 (1992).
- 84 Wei, C., Gershowitz, A., and Moss, B., N6, O2'-dimethyladenosine a novel methylated ribonucleoside next
to the 5' terminal of animal cell and virus mRNAs. *Nature* **257** (5523), 251 (1975); Wei, C. M. and Moss,
B., Methylated nucleotides block 5'-terminus of vaccinia virus messenger RNA. *Proc Natl Acad Sci U S A*
72 (1), 318 (1975).
- 85 Gordon, K. H., Johnson, K. N., and Hanzlik, T. N., The larger genomic RNA of Helicoverpa armigera
stunt tetravirus encodes the viral RNA polymerase and has a novel 3'-terminal tRNA-like structure.
Virology **208** (1), 84 (1995); Karasev, A. V. et al., Complete sequence of the citrus tristeza virus RNA
genome. *Virology* **208** (2), 511 (1995).
- 86 Magden, J. et al., Virus-specific mRNA capping enzyme encoded by hepatitis E virus. *J Virol* **75** (14),
6249 (2001).
- 87 Raju, R. et al., Nontemplated bases at the 5' ends of Tacaribe virus mRNAs. *Virology* **174** (1), 53 (1990);
Morin, B. et al., The N-terminal domain of the arenavirus L protein is an RNA endonuclease essential in
mRNA transcription. *PLoS Pathog* **6** (9).
- 88 Reguera, J., Weber, F., and Cusack, S., Bunyaviridae RNA polymerases (L-protein) have an N-terminal,
influenza-like endonuclease domain, essential for viral cap-dependent transcription. *PLoS Pathog* **6** (9).
- 89 Mao, X. and Shuman, S., Intrinsic RNA (guanine-7) methyltransferase activity of the vaccinia virus
capping enzyme D1 subunit is stimulated by the D12 subunit. Identification of amino acid residues in the
D1 protein required for subunit association and methyl group transfer. *J Biol Chem* **269** (39), 24472
(1994).
- 90 Ho, C. K., Van Etten, J. L., and Shuman, S., Expression and characterization of an RNA capping enzyme
encoded by Chlorella virus PBCV-1. *J Virol* **70** (10), 6658 (1996).